

Presynchronization with CIDR improved the efficiency of ovsynch in the cyclic and acyclic Postpartum Pluriparous Cows

Ahmed E. Ahmed^{1*}, Alaa Eldin Zain Elabdeen² and Omar Zain²

¹Theriogenology Department, Faculty of Veterinary Medicine, South Valley University, Qena, 83523, Egypt

²Theriogenology Department, Faculty of Veterinary Medicine, Sohag University, Sohag, 82524, Egypt

Article History: Received: 6/2/2017

Received in revised form: 10/3/2017

Accepted: 20/3/2017

Abstract

The postpartum period is considered an important period in the reproductive life of dairy cows. This study was aimed to evaluate the effect of progesterone (P_4)-CIDR supplementation in the course of Ovsynch regimens in postpartum Holstein-Friesian dairy cows. Eighty Holstein Friesian dairy cows were used. All animals were gynaecologically examined for uterine involution and ovarian resumption by rectal ultrasound scan. Those with history of endometritis, caesarean section, dystocia, retained placenta, acute mastitis, lameness or other unhealthy were excluded. Fifty-two cows were cyclic and twenty eights were non-cyclic. Animals were divided into three synch-regimens. The 1st group (N=27), used as a control, is the ovsynch regimen only in which the animals were treated with GnRH-PGF_{2a}-GnRH (GPG), the 2nd group (N=26) in which the GPG was presynchronized with controlled internal drug release (CIDR) device before (CIDR-GPG), and the 3rd group (N=27) in which CIDR device was incorporated into GPG, after the first GnRH injection (G-CIDR-PG). Blood samples were collected 5 times (day -7, 0, 7, 14 and 50) for measuring P_4 using enzyme-immunoassay (EIA) technique. Ultrasound examinations were performed on days -7, 0, 7 and 9 for following up the ovarian changes, and repeated at 35, 45 and 50 days post-insemination for pregnancy diagnosis. Conception rate was variable among groups. It was increased by 69.23, 51.85 and 40.74% in CIDR-GPG, G-CIDR-PG regimens treated groups and GPG-control group, respectively. The conception rate was higher in the cyclic versus acyclic GPG group (44.4 vs. 33.3 %). On the other hand, acyclic cows were highly responsive than cyclic cows in CIDR-GPG group (70.0 vs. 68.8%) and G-CIDR-PG group (55.6 vs. 50.0%). In conclusion, we recommend that the CIDR-GPG regimen is improving fertility and conception rate in postpartum dairy cows.

Keywords: CIDR, Cow, Conception, Ovsynch

Introduction

The animal's reproductive performance greatly affects its productivity and subsequently its profitability [1]. The calving interval should average one year [2]. Therefore, the interval between parturition and first service should be less than 90 days [3]. It could be achieved by the early resumption of ovarian activity and higher conception rates. However, most of dairy farms depend on artificial insemination (AI) in reproduction, conception rates were dramatically decreased from 66% to about 35% since 1973 till 2000 parallel to the use of AI and higher dry matter intake [4,5]. Resumption of the ovarian activity is controlled by hormonal and environmental factors. Cows respond to the pulsatile release

of luteinizing hormone (LH) generally exhibited a corpus luteum (CL) and subsequently raise the circulating progesterone (P_4) within the first month after parturition [4]. Many different treatment regimens are applied to synchronize estrus. Mostly, a luteolytic agent prostaglandin-F_{2a} (PGF_{2a}) is included either alone or in combination with a regulatory agent of the follicular development. Depending on the fact that progesterone (P_4) is necessary to fertility and its level before insemination was found to be positively correlated to the subsequent conception rate [6,7].

The alternative regimens such as, P_4 , PGF_{2a} and follicle wave regulating agent were widely applied. Our hypothesis was that P_4

*Corresponding author email: (ezzat_29@yahoo.com), Theriogenology Department, Faculty of Veterinary Medicine, South Valley University, Qena, Egypt, 83523.

enhances a rebound phenomenon of the hypothalamic-pituitary-gonadal (HPG) axis. Using P₄ before the ovsynch protocol, could improve the follicular development, oocyte maturation and conception rate. The controlled-internal drug release (CIDR; a pessary progestin-containing device) was incorporated with the ovsynch protocol and found to improve the conception rates of first-lactation cows [8,9]. The aim of the present study was to evaluate the effect of presynchronization with P₄ before ovsynch or incorporation P₄ with the ovsynch on the conception rate in cyclic and acyclic cows.

Material and methods

Animals and management

This study was conducted on governmental dairy farm in South Egypt (Qena province) containing pure Holstein Friesian dairy cows during the period from May to November (ambient temperature, 34-40 °C and humidity, 60-70%). Animals were fed a total mixed ration according to the National Research Council [10] with continuously available water. After calving, the newborn calves were allowed to suckle their dams for five days. Cows were milked twice daily at 12 h interval. The external signs of estrous were checked twice daily by herdsmen. All animals were bred by natural mating at the first observed estrus (usually 46 days post-partum). Pregnancy was diagnosed by rectal palpation 50-60 days after mating. Eighty healthy pluriparous cows with history of normal calving (age, 4.2 ± 0.1 years old; milk yield average, 6405 ± 0.36 kg/year/cow) were used. The body condition score (BCS) of cows averaged 3.0 ± 0.04 (BCS, 1= emaciated and 5 = obese) [11], the values were presented as mean ± SEM. The experiment started at the 30th days post-partum (P.P) after detecting the cyclicity state of the animals.

Hormonal drugs

Gonadotropin-releasing hormone (GnRH) (Receptal, Intervet Egypt Co. for animal health, 1 mL contains 0.0042 mg buserelin acetate equivalent to 0.004 mg buserelin) [12]. Prostaglandins-F_{2α} (PGF_{2α}) (Lutalyse, Zoetis Animal Health, 1 mL contains 5 mg dinoprost) [13]. The controlled internal drug-release (CIDR) (Pfizer Animal Health,

silicon-coated devices containing 1.38 g of progesterone/device inserted intravaginally) [14].

Experimental design

Cows were classified based on the gynecological examination (rectal palpation for uterine involution), ovarian-ultrasound scan and vaginoscopy (for secretions) into cyclic (N=52) and acyclic (N=28) based on the presence of the corpus luteum (CL) or follicles [15] and the reports of the regular monthly examination for the state of the animals and confirmed by the ultrasonography and hormonal analysis that matching the previous report specific to each animal in the farm. The first ovarian examination by ultrasonography was started on the 20th day P.P and repeated 10 days later for detecting the CL and follicular activity. The start of the study was defined as day -7 and the cows were allocated to one of the following groups as shown in Fig. 1. The first group (GPG) contained twenty-seven animals (cyclic 18 cows and non-cyclic 9 cows) that incorporated into ovsynch protocol (GPG), while the second group (CIDR-GPG) contained twenty six animals (cyclic 16 cows and non-cyclic 10 cows) that incorporated into G-CIDR-PG protocol and the third group (G-CIDR-PG) contained twenty seven animals (cyclic 18 cows and non-cyclic 9 cows) that incorporated into G-CIDR-PG protocol. Blood samples were collected 5 times from jugular vein in heparinized tubes (Day -7, 0, 7, 14 and 50). The harvested plasma samples were stored at -20 °C until the estimation of P₄ concentrations. Plasma P₄ was measured by enzyme-linked immune sorbent assay (ELISA) [16] using progesterone Kits (Immunospec corporation, U.S.A).

Ultrasonographic examinations were carried out on days -7 (start of treatment), 0, 7, 9 and 50 for all animals in order to record the ovarian cyclicity, response to GnRH (Intervet Egypt Co. for animal health, 1ml contains 0.0042 mg buserelin acetate equivalent to 0.004 mg buserelin), luteolytic effect of PGF_{2α} (Zoetis Animal Health, 1 mL contains 5 mg dinoprost), and the pregnancy diagnosis, respectively. Pregnancy diagnosis was recorded by 34 days after mating or 50 days

from the start of the protocol. Cows with detectable luteal tissue on Day -7 were considered cyclic. For recording the conception rate, pregnancy was determined at 35-45 days after mating by using the real time B-mode ultrasound scanner (DRAMINSKI ANIMAL profile scanner equipped with a 7.5 MHz linear transducer; DRAMINSKI Co., Poland).

Statistical analysis

Chi-square was used for analyzing the data of conception rates. All data of plasma P₄

concentrations were tested for significant difference by one-way ANOVA and confirmed by Newman-Keuls as a post-hoc test. All data of milk yield between cyclic and acyclic animals in the different groups of the study were tested for significant difference by two-way ANOVA and confirmed by Bonferroni as a post-hoc test. All data were presented as mean ± S.E.M. The differences were considered significant at P<0.05 using Graph Pad Prism software program (V.5.01, San Diego USA, 2007).

Table 1: Descriptive statistics of body condition score (BCS), parity and milk production (kg/day) in postpartum dairy cows at the start of treatments; ovsynch protocol (GPG), presynchronization of ovsynch with CIDR (CIDR-GPG) and incorporation on CIDR into the ovsynch (G-CIDR-PG). All values were presented as Mean ± S.E.M

| Treatment | Number | B.C.S. | Parity |
|------------------------|-----------|--------------------|-------------------|
| GPG group | | | |
| Non-cyclic | 9 | 2.27 ± 0.12 | 1.44 ± 0.17 |
| Cyclic | 18 | 2.75 ± 0.08 | 1.22 ± 0.10 |
| Total | 27 | 2.59 ± 0.08 | 1.30 ± 0.09 |
| CIDR-GPG group | | | |
| Non-cyclic | 10 | 3.25 ± 0.83 | 1.70 ± 0.15 |
| Cyclic | 16 | 3.34 ± 0.08 | 1.94 ± 0.06 |
| Total | 26 | 3.31 ± 0.06 | 1.85 ± 0.07 |
| G-CIDR-PG group | | | |
| Non-cyclic | 9 | 3.22 ± 0.87 | 2.0 ± 0.00 |
| Cyclic | 18 | 3.25 ± 0.06 | 2.0 ± 0.08 |
| Total | 27 | 3.24 ± 0.05 | 2.0 ± 0.05 |

Results

The descriptive statistics of BCS, parity and milk production (kg/day/cow) of animals in the three groups; GnRH-PGF_{2α}-GnRH (GPG), controlled-internal-drug-release (CIDR)-GPG and G-CIDR-PG groups in cyclic and acyclic cows were shown in table 1. No significant difference was recorded for milk production between the cyclic and acyclic cows within each protocol of study. Conception rates after natural mating following GPG, CIDR-GPG and G-CIDR-PG in cyclic and acyclic postpartum dairy cows were shown in table 2. The conception rate tended to be greater in CIDR-based protocols than in GPG-ovsynch protocol. Furthermore, conception rate varied in response to different groups. In GPG group, the acyclic animals showed lower conception rates than those cyclic (33.3 vs. 44.4 %) but the response was reversed in the CIDR-treated ovsynch groups where the acyclic animals showed higher

conception rates compared to cyclic in the CIDR-GPG (70.0 vs. 68.8 %) and G-CIDR-PG (55.6 vs. 50.0 %) (Table 2). Regarding the milk yield, there was no significant difference between cyclic and acyclic cows among the different groups of the study.

Figure 2 shows the ultrasound images of ovarian structures, including; the follicle (Fig. 2A) and corpus luteum (Fig. 2B) on days -7, 0, 7, 9 and 14, in addition to the amniotic vesicle (Fig. 2C) as a sign of pregnancy diagnosis was recorded on day 34 after mating that equals the day 50 from the start of the protocol. Plasma P₄ concentrations in cyclic and acyclic cows after three protocols of estrus synchronization were shown in Fig. 3. The data related to pregnant cyclic and acyclic animals were presented in Fig. 3A, and 3C, respectively. The plasma P₄ levels on day 0 were significantly higher (P<0.05) in CIDR-GPG protocol compared to other

treatment groups. However, its levels on day 7 were significantly higher ($P<0.05$) in G-CIDR-PG protocol compared to other treatment groups. For the non-pregnant cyclic and acyclic animals (Fig. 3B, and 3D, respectively), the plasma P₄ profile from day 0 up to day 7 revealed the same results as mentioned for those cyclic. The cyclic

animals, both pregnant and non-pregnant, the plasma P₄ levels on day 14 maintained significantly higher in CIDR protocols rather than non-CIDR ovsynch regimen ($P<0.05$) (Fig. 3A and 3B). However, the acyclic animals showed non-significant differences in response to all ovsynch regimens on the same day (Fig. 3C and 3D).

Table 2: Conception rates after natural mating following GPG, CIDR-GPG and G-CIDR-PG in cyclic and acyclic postpartum dairy cows

| Experimental groups | Cyclic | | | | Acyclic | | | | Conception rate |
|---------------------|--------|----------|----|--------------|---------|----------|----|--------------|-----------------|
| | N | Pregnant | N | Non-pregnant | N | Pregnant | N | Non-pregnant | |
| GPG | 8 | 44.4% | 10 | 55.6 % | 3 | 33.3 % | 6 | 66.7 % | 40.74 % |
| CIDR-GPG | 11 | 68.8% | 5 | 31.25% | 7 | 70.0 % | 3 | 30.0 % | 69.23 % |
| G-CIDR-PG | 9 | 50.0% | 9 | 50.0% | 5 | 55.6 % | 10 | 44.4 % | 51.85 % |

N: Number; Chi square=8.33, df = 1, P = 0.0039**

Discussion

Several modifications of ovsynch were applied depending on hormonal supplementation of progestin, PGF_{2α} and gonadotropins [17,18]. One of the causes of poor fertility in high producing dairy cows is inadequate progesterone and 60-85% of dairy cows showed a suboptimal circulating P₄ for pregnancy [19]. On dependence on evaluating the effect of P₄ supplementation throughout

the course of Ovsynch protocol in cyclic and acyclic P.P Holstein Friesian dairy cows, we recorded an increase in the conception rate when P₄ was used with the Ovsynch protocol. This result was in accordance with Kawate *et al.* [20] who reported that the addition of CIDR to the ovsynch protocol (G-CIDR-PG) improved the conception rates in postpartum Japanese Black beef cows.




Figure 1: Experimental design for the studied groups: (A) GPG group: Day 0, GnRH (Receptal, 5 mL intramuscular injection); Day 7, PGF_{2α} (Lutalyse, 5 mL IM); Day 9, GnRH was repeated followed by natural mating after estrous detection that monitored 16-18 h after the last GnRH. (B) CIDR-GPG: Day -7, cows were initially treated with CIDR intravaginally for 7 days; Day 0, GnRH (Receptal, 5 mL IM) and removal of CIDR's devices; Day 7, PGF_{2α} (Lutalyse, 5 mL IM); Day 9, GnRH was repeated followed by natural mating after estrous detection that monitored 16-18 h after the last GnRH. (C) G-CIDR-PG: Day 0, GnRH (Receptal, 5 mL IM) and insertion of CIDR's devices for 7 days; Day 7, PGF_{2α} (Lutalyse, 5 mL IM) and removal of CIDR's devices; Day 9, GnRH was repeated followed by natural mating after estrous detection that monitored 16-18 h after the last GnRH.

The current investigation presented presynchronization with CIDR before GPG as a new regimen as an attempt to enhance the conception rate in comparison to ovsynch or G-CIDR-PG. Conception rate was variable among groups. It is likely that the addition of P₄ to the Ovsynch protocol may be affected by a number of variables such as age, post-partum interval and ovarian follicular growth. Pregnancy rates were positively associated with higher P₄ concentration in the luteal phase of the cycle preceding AI [21]. This proved by the findings that reported an

improvement in the endometrial morphology following elevated P₄ concentrations of the preceding cycle. The high pregnancy rates were attributed to the quadratic effect of P₄ at the PGF_{2α} injection in ovsynch protocol. Cows with P₄ ≥ 1 ng/mL at the PGF_{2α} injection had greater pregnancy rate (41%) than cows with P₄ ≤ 1 ng/mL (12%) [22]. Variations in serum P₄ at the PGF_{2α} injection dramatically affected fertility in dairy cows. Higher P₄ concentrations prior to insemination reduced uterine secretion of PGF_{2α} (in response to oxytocin) during late luteal phase [23].



Figure 2: Representative ultrasound image of ovarian activity and follicular dynamics in synchronized cows of all groups at start of treatment (day -7) showing growing follicle (A), corpus luteum (B). Ultrasound image of uterine horn in a studied cow shows the embryonic vesicle at around day 40 post mating (C).

In addition, high P₄ levels at the time of PGF_{2α} injection and luteal regression shortly after PGF_{2α} treatment induced a greater response to GnRH and better fertility in lactating dairy cows [24]. Lamb *et al.* [25] reported that P₄-CIDR supplementation between GnRH and PGF_{2α} treatment could improve the pregnancy rate in cows with low P₄ concentrations at PGF_{2α} injection. Progesterone was considered a ‘proxy variable’ among other factors associated with conception rate. The fact that ovulation induced by GnRH administration followed by a high proportion of pre-mature luteal regression can be avoided by priming with exogenous P₄ [26]. Our results were incorporated with other studies that reported CIDR insertion improved the conception rates in non-cyclic cows and the exogenous

supplement of P₄ served as a primer to facilitate ovulation of a follicle in response to the second GnRH injection [25-27]. The conception rates were higher in cyclic cows than in acyclic cows, which agree with Geary *et al.* [28] but contradict the report of Sakase *et al.* [29] who stated no effect of cyclicity on the conception rates in response to ovsynch.

In the present study, concentrations of P₄ in cows with or without an active CL were increased by the CIDR insertion and P₄ increased in both CIDR-GPG and G-CIDR-PG on day 0 and 7, respectively. These elevations are in agreement with those achieved by different authors [20,30,31] who indicated that P₄ concentrations were rapidly increased and reached its peak within 1h after CIDR insertion and then decreased rapidly up to 0 ng/mL 12 to 24 h once the CIDR was

removed. Other studies concluded that CIDR insertion provided a relatively small increase (0.5 to 1 ng/mL) in serum P₄ concentrations [27]. The CIDR might maintain P₄ above 1 ng/mL for at least 7 days in cows without CL [28]. The increase in P₄ from day 0 to 7 was small in the ovsynch group. In cycling cows, plasma P₄ concentration tended to be higher

on days 0 and 7 in the CIDR-GPG and G-CIDR-PG protocols than those in the ovsynch protocol. Moreover, in GPG protocol, the first GnRH treatment was given on approximately day 15 of the estrus cycle and the CL was regressed before the PGF_{2α} treatment and estrus occurred, resulting in ovulation before timed-AI [32].




Figure 3: Plasma P₄ concentrations of pregnant cyclic (A), non-pregnant cyclic (B), pregnant acyclic (C), and non-pregnant acyclic (D) cows in response to treatment protocols of GPG, CIDR-GPG and G-CIDR-PG groups throughout the periods of study. The letters on bars (a, b and c) denote the significant difference at *P<0.05. NS means non-significant difference.

In acyclic cows, plasma P₄ concentrations were clearly higher on day 0 in CIDR-GPG group than in the ovsynch group that could be due to the release of P₄ from CIDR. On day 7, plasma P₄ concentrations were clearly higher in G-CIDR-PG group than ovsynch and CIDR-GPG groups that may be related to the release of P₄ from CIDR, in addition to a possible ovulation or luteinization following the first GnRH treatment. The plasma P₄ concentration was (8.81 ± 0.71 ng/mL) at day 50 in all conceived cyclic cows, this result came in agreement with that obtained by Han *et al.* [33] who found that P₄ at 15 to 32 days after AI was

consistently higher in pregnant (> 4 ng/mL) than non-pregnant cows.

In non-pregnant cows, P₄ concentrations in CIDR-GPG and G-CIDR-PG groups increased on days 0 and 7 than in the ovsynch protocol, which attributed to the release of P₄ from CIDR. The P₄ concentrations on day 14 were high (3 up to 6 ng/mL), which referred to the efficiency of the ovsynch protocols (with or without CIDR) in stimulating the resumption of ovarian activity and regardless the incidence of conception in both cyclic and acyclic groups. The variations of P₄ concentration among groups is related to the follicular size at ovulation. Induction of ovulation in cows with smaller dominant

follicle at the second GnRH treatment resulted in smaller CL [34] and therefore, reduced the fertility and P₄ concentration [35].

In the pregnant cows, the P₄ concentration after insemination was lower in cows of non-CIDR ovsynch protocol than those subjected to CIDR and that conception rate varies among groups recording the lowest values in the non-CIDR's animals. Sartori *et al.* [36] and Wolfenson *et al.* [37] suggested that prolonged exposure to P₄ increase the rhythm of the follicular growth leading to larger follicles and consequently larger corpora lutea that produce more P₄ [38]. In addition, P₄ level is the most important factor affecting embryonic life in dairy cows. Previous studies showed that embryonic loss was high in cows with low levels of P₄ compared to those of high levels during the first week after insemination [39].

The results of the current study revealed that the overall conception rate in GPG and G-CIDR-PG groups were 40.74% and 51.85%, respectively that approximately similar to the results reported by Stevenson *et al.* [40] who stated that conception rates at the day 28th post insemination was 50% following ovsynch plus CIDR. On the other hand, presynchronization with CIDR before ovsynch protocol was improved the conception rate to 69.23 %. Prolonged effect of P₄ (CIDR and induced CL) lead to delay in LH release, which allow smaller follicles to fully mature and increased the ova number [41]. From the genetic view, prolonged maintenance of oocyte in the follicular environment could be beneficial to the accumulation of more maternal mRNA as a good factor for the embryonic growth [41, 42].

Conclusion

The current study presented that the CIDR-GPG protocol as a new ovsynch modification could be used for estrus synchronization in the postpartum dairy cows. It achieved a higher conception rate especially in acyclic cows, while the conception rates in all treatments was affected by the cyclic status of animals.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

Our deepest thanks to Dr. Mohammed Emad Farrag and Dr. Ahmed Abd El-hallem Waked, (Large Animal, Zoetis, Middle East), for supplying the laboratory with: CIDR and Lutalyse.

References

- [1] Britt, J.H. (1975): Early postpartum breeding in dairy cows. A review1. *J Dairy Sci*, 58(2): 266-271.
- [2] Louca, A. and Legates, J.E. (1968): Production losses in dairy cattle due to days open. *J Dairy Sci*, 51(4): 573-583.
- [3] Fagan, J.G. and Roche, J.F. (1986): Reproductive activity in postpartum dairy cows based on progesterone concentrations in milk or rectal examination. *Irish Vet J*, 40(7-8): 124-131.
- [4] Butler, W.R. and Smith, R.D. (1989): Interrelationships between energy balance and postpartum reproductive function in dairy cattle. *J Dairy Sci*, 72(3): 767-748.
- [5] Lucy, M.C. (2001): Reproductive loss in high-producing dairy cattle: Where will it end? *J Dairy Sci*, 84(6):1277-1293.
- [6] Fonseca, F.A.; Britt, J.H.; McDaniel, B.T.; Wilk, J.C. and Rakes, A.H. (1983): Reproductive traits of Holsteins and Jerseys. Effects of age, milk yield, and clinical abnormalities on involution of cervix and uterus, ovulation, estrous cycles, detection of estrus, conception rates and days open. *J Dairy Sci*, 66(5):1128-1147.
- [7] Folman, Y.; Kaim, M.; Herz, Z. and Rosenberg, M. (1990): Comparison of methods for the synchronization of estrous cycles in dairy cows. 2. Effects of progesterone and parity on conception. *J Dairy Sci*, 73(10): 2817-2825.
- [8] El-Zarkouny, S.Z.; Cartmill, J.A.; Hensley, B.A. and Stevenson, J.S. (2004): Pregnancy in dairy cows after synchronized ovulation regimens with or without presynchronization and progesterone. *J Dairy Sci*, 87(4): 1024-1037.
- [9] Moreira, F.; Flores, R. and Boucher, J. (2004): Use of CIDR with a timed insemination protocol in lactating dairy

- cows during summer in Mexico. *J Dairy Sci*, 87: 373.(Abstract).
- [10] NRC. (1989): Nutrient Requirements of Dairy Cattle. 5th rev. ed. Natl. Acad. Sci. Washington, DC.
- [11] Spitzer, J.C. (1986): Influences of nutrition on reproduction in beef cattle. In Morrow, D.A. Current therapy in Theriogenology 2nd ED. W.B. Sounders, Philadelphia PA, p:320-340.
- [12] Nawito, M., Schallenberger, E. and Schams, D. (1977): Release of lutropin (LH) and follitropin (FSH) in cattle after administration of a new gonadoliberin (GnRH) analogue in comparison with the gonadoleberin decapeptide. *Theriogenology*, 7: 277-291.
- [13] Moody, E.L. (1979): Studies on Lutalyse use programs for estrus control. Proceedings of the Lutalyse Symposium, Brook Lodge, Augusta, MI, p.33-41.
- [14] Rathbone, M.J.; Bunt, C.R.; Ogle, C.R.; Burggraaf, S.; Macmillan, K.L.; Burke, C.R. and Pickering, K.L. (2002): "Reengineering of a commercially available bovine intravaginal insert (CIDR insert) containing progesterone". *Journal of Controlled Release*, 85: 105-115.
- [15] De Rensis, F.; Ronci, G.; Guarneri, P.; Nguyen, B.X.; Presicce, G.A.; Huszenicza, G.Y. and Scaramuzzi, R.J. (2005): Conception rate after fixed time insemination following ovsynch protocol with and without progesterone supplementation in cyclic and non-cyclic Mediterranean Italian buffaloes (*Bubalus bubalis*). *Theriogenology*, 63(7): 1824-1831.
- [16] Elder, P.A.; Yeo, K.H.J.; Lewis J.G. and Clifford, J.K. (1987): An Enzyme-linked immunosorbent assay (ELISA) for plasma progesterone: immobilized antigen approach. *Clin Chim Acta*, 162(2): 199-206.
- [17] Odde, K.G. and Holland, M.D. (1994): Synchronization of estrus in cattle. In: Factors Affecting Calf Crop. CRC Press, Boca Raton, FL.
- [18] Ryan, D.P.; Snijders, S.; Yaakub, H. and O'Farrell, K.J. (1995): An evaluation of estrus synchronization programs in reproductive management of dairy herds. *J Anim Sci*, 73(12): 3687-3695.
- [19] Stronge, A.J.H.; Sreenan, J.M.; Diskin, M.G.; Mee, J.F.; Kenny, D.A. and Morris, D.G. (2005): Post-insemination milk progesterone concentration and embryo survival in dairy cows. *Theriogenology*, 64(5):1212-1224.
- [20] Kawate, N.; Itami, T.; Chousi, T.; Saitoh, T.; Wada, T. and Matsuoka, K. (2004): Improved conception in timed-artificial insemination using a progesterone-releasing intravaginal device and Ovsynch protocol in postpartum suckled Japanese Black beef cows. *Theriogenology*, 61(12): 399-406.
- [21] Rosenberg, M.; Kaim, M.; Herz, Z. and Folman, Y. (1990): Comparison of methods for synchronization of estrous cycles in dairy cows. Effects on plasma progesterone and manifestation of oestrus. *J Dairy Sci*, 73(10): 2807-2816.
- [22] Sterry, R.A.; Silva, E.; Kolb, D. and Fricke, P.M. (2009): Strategic treatment of anovular dairy cows with GnRH. *Theriogenology*, 71(3): 534-542.
- [23] Shaham-Albalancy, A.; Folman, Y.; Kaim, M.; Rosenberg, M. and Wolfenson, D. (2001): Delayed effect of low progesterone concentrations on bovine uterine PGF2a secretion in the subsequent oestrous cycle. *Reprod*, 122(4): 643-648.
- [24] Moreira, F.; Orlandi, C.; Risco, C.A.; Mattos, R.; Lopes, F. and Thatcher, W.W. (2001): Effects of presynchronization and bovine somatotropin on pregnancy rates to timed artificial insemination protocol in lactating dairy cows. *J Dairy Sci*, 84(7): 1646-1659.
- [25] Lamb, G.C.; Stevenson, J.S.; Kesler, D.J.; Garverick, H.A.; Brown, D.R. and Salfen, B.E. (2001): Inclusion of an intravaginal progesterone insert plus GnRH and prostaglandin F2 for ovulation

- control in postpartum suckled beef cows. *J Anim Sci*, 79(9): 2253-2259.
- [26] Larson, J. E.; Lamb, G.C.; Stevenson, J.S.; Johnson, S.K.; Day, M.L.; Geary, T.W.; Kesler, D.J.; De Jarnette, J.M.; Schrick, F.N.; DiCostanzo, A. and Arseneau, J.D. (2006): Synchronization of estrus in suckled beef cows for detected estrus and artificial insemination and timed artificial insemination using gonadotropin-releasing hormone, prostaglandin F_{2α}, and progesterone. *J Anim Sci*, 84(2): 332-342.
- [27] Stevenson, J.S.; Lamb, G.C.; Johnson, S.K.; Medina-Britos, M.A.; Grieger, D.M.; Harmoney, K.R.; Cartmill, J.A.; El-Zarkouny, S.Z.; Dahlen, C.R. and Marple, T.J. (2003): Supplemental norgestomet, progesterone, or melengestrol acetate increases pregnancy rates in suckled beef cows after timed inseminations. *J Anim Sci*, 81(3): 571-586.
- [28] Geary, T.W.; Whittier, J.C.; Hallford, D.M. and MacNeil, M.D. (2001): Calf removal improves conception rates to the Ovsynch and CO-Synch protocols. *J Anim Sci*, 79(1): 1-4.
- [29] Sakase, M.; Seo, Y.; Fukushima, M.; Noda, M.; Takeda, K.; Ueno, S.; Inaba, T.; Tamada, H.; Sawada, T. and Kawate, N. (2005): Effect of CIDR-based protocols for timed-AI on the conception rate and ovarian functions of Japanese Black beef cows in the early postpartum period. *Theriogenology*, 64(5): 1197-1211.
- [30] Perry, G.A.; Smith, M.F. and Geary, T.W. (2004): Ability of intravaginal progesterone inserts and melengestrol acetate to induce estrous cycles in postpartum beef cows. *J Anim Sci*, 82(3): 695-704.
- [31] Lamb, G.C.; Larson, J.E.; Geary, T.W.; Stevenson, J.S.; Johnson, S.K.; Day, M.L.; Ansotegui, R.P.; Kesler, D.J.; DeJarnette, J.M. and Landblom, D.G. (2006): Synchronization of estrus and artificial insemination in replacement beef heifers using gonadotropin-releasing hormone, prostaglandin F_{2α}, and progesterone. *J Anim Sci*, 84(11): 3000-3009.
- [32] Moreira, F.; De la Sota, R.L.; Diaz, T. and Thatcher, W.W. (2000): Effect of day of the estrous cycle at the initiation of a timed artificial insemination protocol on reproductive responses in dairy heifers. *J Anim Sci*, 78(6): 1568-1576.
- [33] Han, H.; Austin, K.J.; Rempel, L.A. and Hansen, T.R. (2006): Low blood ISG15 mRNA and progesterone levels are predictive of non-pregnant dairy cows. *J Endocrinol*, 191(2): 505-512.
- [34] Macmillan, K.L.; Segwagwe, B.V. and Pino, C.S. (2003): Associations between the manipulation of patterns of follicular development and fertility in cattle. *Anim Reprod Sci*, 78(3): 327-344.
- [35] Cavalieri, J.; Hepworth, G. and Macmillan, K.L. (2004): Ovarian follicular development in Holstein cows following synchronization of oestrus with oestradiol benzoate and an intravaginal progesterone releasing insert for 5-9 days and duration of the oestrous cycle and concentrations of progesterone following ovulation. *Anim Reprod Sci*, 81(3): 177-193.
- [36] Sartori, R.; Sartor-Bergfelt, R.; Mertens, S.A.; Guenther, J.N.; Parrish, J.J. and Wiltbank, M.C. (2002): Fertilization and early embryonic development in heifers and lactating cows in summer and lactating and dry cows in winter. *J Dairy Sci*, 85(11): 2803-2812.
- [37] Wolfenson, D.; Inbar, G.; Roth, Z.; Kaim, M.; Bloch, A. and Braw-Tal, R. (2004): Follicular dynamics and concentrations of steroids and gonadotropins in lactating cows and nulliparous heifers. *Theriogenology*, 62(6): 1042-1055.
- [38] Perry, G.A.; Smith, M.F.; Lucy, M.C.; Green, J.A.; Parks, T.E.; MacNeil, M.D.; Roberts, A.J. and Geary, T.W. (2005): Relationship between follicle size at insemination and pregnancy success. *Proc Natl Acad Sci USA*, 102: 5268-5273.
- [39] Lopez-Gatius, F.; Santolaria, P.; Martino, A.; Deletang, F. and De Rensis, F.

- (2006): The effects of GnRH treatment at the time of AI an 12 days later on reproductive performance of high producing dairy cows during the warm season in northeastern Spain. *Theriogenology*, 65(4): 820-830.
- [40] Stevenson, J.S.; Pursley, J.R.; Garverick, H.A.; Fricke, P.M.; Kesler, D.J.; Ottobre, J.S. and Wiltbank, M.C. (2006): Treatment of cycling and noncycling lactating dairy cows with progesterone during Ovsynch. *J Dairy Sci*, 89(7): 2567-2578.
- [41] Merton, J.S.; De Roos, A.P.; Mullaart, E.; De Ruigh, L.; Kaal, L.; Vos, P.L. and Dieleman, S.J. (2003): Factors affecting oocyte quality and quantity in commercial application of embryo technologies in the cattle breeding industry. *Theriogenology*, 59(2): 651-674.
- [42] Vassena, R.; Mapletoft, R.J.; Allodi, S.; Singh, J. and Adams, G.P. (2003): Morphology and developmental competence of bovine oocytes relative to follicular status. *Theriogenology*, 60(5): 923-932.

الملخص العربي

تحسين معدل حدوث الحمل والخصوبة باستخدام السيدار قبل نظام تزامن التبويض في فترة ما بعد الولادة بالأبقار الحلاة
أحمد عزت أحمد^١ ، علاء الدين زين العابدين محمود^٢ ، عمر زين العابدين^٣

^١قسم التناهيات والتوليد والتلقيح الاصطناعى بكلية الطب البيطري جامعة جنوب الوادى. ص.ب ٨٣٥٢٣ قنا - جمهورية مصر العربية

^٢قسم التناهيات والتوليد والتلقيح الاصطناعى بكلية الطب البيطري جامعة وهاج. ص.ب ٨٢٥٢٤ وهاج - جمهورية مصر العربية

تعتبر فترة ما بعد الولادة مهمة جداً للحياة التالية للأبقار الحلاة. تهدف الدرة الحالية إلى تحدى البروجيسترون (P4) والسيدار المهبلي في برنامج تزامن دورة التبويض والشيق في أبقار الهولشتاين فريزيان الحلاة. تناولت الدرة عدد ٨٠ بقرة حلاة. جميع الحيوانات تم فحصها تناهياً من حيث عودة الجهاز التناسلي مثل الرحم والتغاير نشاط المبيض وذلك من خلال الجس والفحص بروطة السونار من خلال فتحة المستقيم. وبعد اجراء الفحوصات الاكلينيكية والاطلاع على السجلات الصحية بالمزرعة يتم تبعيد الحيوانات التي تعاني من التهابات رحمية أو مهبليّة أو يبقى أن تعرّضت لعسر ولادة وعمليات قصيرة واحتباس المშيمة والعرج والتهاب الضرع. اثنان وخمسون حيواناً منتظماً في دورة الشيق بينما ثمانية وعشرون حيواناً غير منتظم في دورة الشيق. تم تقسيم الحيوانات إلى ثلاثة مجموعات حسب تأثيرهم المعاملات الدرية إلى الآتي:- المجموعة الأولى وعدها ٢٧ حيوان تلقت فقط النظرة العادي أو جي-بي-جي بينما المجموعة الثانية وعدها ٢٦ حيوان تم معاملتها بالسيدار المهبلي قبل نظر الجي-بي-جي، أما المجموعة الثالثة وعدها ٢٧ حيوان تم معاملتها بالسيدار المهبلي بعد معاملة الحيوانات بالمحفز التناهيلي GnRH لنظرة الجي-بي-جي. تمأخذ عينات كل خمس مرات في اليوم السابع قبل المعاملات - اليوم صفر لبدء المعاملات - واليوم السادس والرابع عشر والخمسين لقياس البروجيسترون. ثم تم الفحص بروطة السونار في اليوم السابع قبل المعاملات - اليوم صفر لبدء المعاملات - واليوم السادس والتاسع ثم إعادة الفحص بالأيام الخامس والثلاثين والخامس والأربعين لتأكيد الحمل. أظهرت النتائج تنوّع معدل الحمل بين المجموعات حيث زاد بنسبة ٦٩.٢٣٪، ٨٥٪، ٥١٪، ٧٤٪ و ٤٠٪ في كل من المجموعة المعاملة الثانية والثالثة مقارنة بالمجموعة الضابطة الأولى على التوالي. وكان معدل الحمل في المجموعة النشيطة أعلى من المجموعة الخامدة في الحيوانات التي تلقت نظرة جي-بي-جي (٤٤.٤٪ مقابل ٣٣.٣٪ على التوالي). بينما تجاوزت الحيوانات الخامدة أكثر من النشيطة في المجموعة الثانية (٧٠٪ مقابل ٦٨.٨٪ على التوالي) وكذلك في المجموعة الثالثة (٥٥.٦٪ مقابل ٥٠٪ على التوالي). بناءً على ما يُذكر فإن نتائج البحث توصي بتناول السيدار المهبلي قبل نظر الجي-بي-جي في تحسين الخصوبة ومعدل الحمل في الأبقار الحلاة في فترة ما بعد الولادة.